

Made in United States of America
Reprinted from THE AUK
Vol. 105, No. 3, July 1988
Copyright © 1988 by The American Ornithologists' Union

Polygyny in the Northern Shrike (*Lanius excubitor*) in Israel

REUVEN YOSEF AND BERRY PINSHOW

*Mirani Center for Desert Ecology, Jacob Blaustein Institute for Desert Research, and
Biology Department, Ben-Gurion University, Sede Boqer Campus 84993, Israel*

Polygyny is the practice of a male having two or more mates, either simultaneously (harem polygyny) or sequentially (successive polygyny) (Lack 1968). Polygyny has been reported in only one species of true shrike (family Laniidae), the Loggerhead (*Lanius ludovicianus*; Verner and Willson 1969 and references therein), despite extensive studies of breeding behavior in several other species (e.g. Bent 1965, Huhtala et al. 1977, Bassin 1982, Kridelbaugh 1983, Dittami and Knauer 1986, Zack 1986).

In the course of a study of Northern Shrikes (*Lanius excubitor*) at Sede Boqer in the Negev Desert, Israel, we observed two cases of harem polygyny. During

the breeding season of 1987 (January-June) 7 pairs of shrikes were observed closely and all individuals were color-banded. Territory boundaries were marked by coordination on a topographical survey map, and areas were measured by planimetry (Table 1).

In Case 1 a pair (male: green band, female: red band) of Northern Shrikes completed building its first nest by 16 January. Five eggs were laid between 26 and 30 January. Incubation by the female began with the laying of the third egg, and 4 nestlings hatched on 11-12 February; two nestlings fledged. On 3 February we observed that a female (blue band), in a territory adjacent to that of the red-banded female, finished

TABLE 1. Sizes of adjacent breeding territories and distances between nests of female Northern Shrikes. Each pair of territories was within that of their polygynous mate. Colors refer to leg bands.

Male	Female	Territory size (ha)	Distance between nests (m)
Green	Red	760	630
	Blue	712	
White	Yellow	788	600
	Black	712	

building her nest. She laid 5 eggs between 17 and 25 February and began incubating after the third egg was laid. We observed that the green-banded male fed the blue-banded female while simultaneously providing food to the red-banded female, which she fed to her nestlings. The nest of the blue-banded female was subsequently destroyed by a storm, and we saw the green-banded male chasing her over the next 3 days until she disappeared.

The green-banded male and the red-banded female prepared a second nest, which was ready by 25 February, and laying commenced on 26 February. This clutch comprised 6 eggs laid at 24-h intervals, and again incubation by the female began with the laying of the third egg.

Case 2 involved a white color-banded male and two females, one with a yellow and the other a black leg band. Initially the white-banded male paired with the yellow-banded female, and their nest was ready on 5 February. She laid eggs from 7 to 12 February. The nestlings hatched on 26 February, and on the same day the nest of the black-banded female in the neighboring territory was completed. The black-banded female laid eggs from 8 March and started incubating from 10 March, after laying the third egg. At this stage we observed the white-banded male feeding the black-banded female on her nest and bringing food to the yellow-banded female, who fed her nestlings. All nestlings in both nests fledged.

In neither case were the females seen to confront

each other along their territorial boundaries. Females alone incubated eggs and brooded young. Neither polygynous male was seen feeding young directly; they both brought food to their mates, who did the feeding. During 109 h of observation of monogamous pairs, however, we saw 17 instances of males feeding young directly.

We thank Joel Brown, Yael Lubin, and Uriel Safriel for their comments on drafts of this manuscript and Steve Zack for his constructive review. This paper is contribution No. 67 of the Mitrani Center for Desert Ecology.

LITERATURE CITED

- BASSIN, P. 1982. Sites de nidification et reproduction de la Pie-grieche Grise, *Lanius excubitor*, dans le nord-ouest de la Suisse (Ajoie, canton du Jura). Nos Oiseaux 36: 333-353.
- BENT, A. C. 1965. Life histories of North American wagtails, shrikes, vireos and their allies. New York, Dover Publ.
- DITTAMI, J. P., & B. KNAUER. 1986. Seasonal organization of breeding and moulting in the Fiscal Shrike (*Lanius collaris*). J. Ornithol. 127: 79-84.
- HUHTALA, K., J. ITAMIES, & H. MIKKOLA. 1977. Beitrag zur Brutbiologie und Ernährung des Raubwurgers (*Lanius excubitor*) im Osterbotten, Finland. Beitr. Vogelkd. 23: 129-146.
- KRIDELBAUGH, A. 1983. Nesting ecology of the Loggerhead Shrike in central Missouri. Wilson Bull. 95: 303-308.
- LACK, D. 1968. Ecological adaptations for breeding in birds. London, Chapman and Hall.
- VERNER, J., & M. F. WILLSON. 1969. Mating systems, sexual dimorphism, and the role of male North American passerine birds in the nesting cycle. Ornithol. Monogr. No. 9.
- ZACK, S. 1986. Behaviour and breeding biology of the cooperatively breeding Grey-backed Fiscal Shrike, *Lanius excubitorius*, in Kenya. Ibis 128: 214-233.

Received 15 September 1987, accepted 19 February 1988.